

Week 5: Quiz

Due

Points 10

Questions 10

Time Limit None

Instructions

There are 10 questions worth 1 point each. You will have 10 minutes to complete the quiz.

Attempt History

	Attempt	Time	Score
LATEST	Attempt 1	33 minutes	10 out of 10

Score for this quiz: **10** out of 10 Submitted Apr 28 at 1:28am
This attempt took 33 minutes.



Correct!

Question 1	1 / 1 pts
CO1 Nausea is difficult to discern in a young child. What question should the clinician ask to determine if a child has nausea?	
<hr/>	
<input checked="" type="radio"/> Are you hungry?	
<hr/>	
<input type="radio"/> Are you sick to your tummy?	
<hr/>	
<input type="radio"/> Are you eating the way you normally eat?	
<hr/>	
<input type="radio"/> Are you nauseous?	

Question 2	1 / 1 pts
<hr/>	
<hr/>	

CO6 A 52-year-old female is suspected of having a gastric ulcer and will undergo an Esophagogastroduodenoscopy (EGD). She is concerned because she heard that gastric ulcers can be malignant. How should the clinician respond?

Correct!

- About 95% of gastric ulcers are benign.
- Don't worry. Gastric ulcers are not cancerous.
- You have about a 50% chance of having gastric cancer from your ulcer.
- Even if it is cancer, surgery is 100% successful.

Question 3

1 / 1 pts

CO6 A 72-year-old male patient presents to the clinic for his annual physical. He states that he is worried because he only has a bowel movement every three days. How should the clinician respond?

Correct!

- There is no such thing as a normal pattern of defecation.
- You should have at least three stools per week.
- You should defecate once a
- You should have two to three stools per